

SECANT ZETA FUNCTIONS

MATILDE LALÍN, FRANCIS RODRIGUE, AND MATHEW ROGERS

ABSTRACT. We study the series $\psi_s(z) := \sum_{n=1}^{\infty} \sec(n\pi z)n^{-s}$, and prove that it converges under mild restrictions on z and s . The function possesses a modular transformation property, which allows us to evaluate $\psi_s(z)$ explicitly at certain quadratic irrational values of z . This supports our conjecture that $\pi^{-k}\psi_k(\sqrt{j}) \in \mathbb{Q}$ whenever k and j are positive integers with k even. We conclude with some speculations on Bernoulli numbers.

1. INTRODUCTION

Let $\zeta(s)$ denote the Riemann zeta function. It is well known that $\zeta(2n)\pi^{-2n} \in \mathbb{Q}$ for $n \geq 1$. Dirichlet L -functions and Clausen functions are modified versions of the Riemann zeta function, which also have nice properties at integer points [Lew91]. Berndt studied another interesting modification of the Riemann zeta function, namely the cotangent zeta function [Ber76]:

$$c_s(z) := \sum_{n=1}^{\infty} \frac{\cot(\pi n z)}{n^s}. \quad (1.1)$$

He proved that (1.1) converges under mild restrictions on z and s , and he produced many explicit formulas for $c_k(z)$, when z is a quadratic irrational, and $k \geq 3$ is an odd integer. Consider the following examples:

$$c_3\left(\frac{1+\sqrt{5}}{2}\right) = -\frac{\pi^3}{45\sqrt{5}}, \quad c_5(\sqrt{2}) = \frac{\pi^5}{945\sqrt{2}}.$$

Berndt's work implies that $\sqrt{j} c_k(\sqrt{j})\pi^{-k} \in \mathbb{Q}$ whenever j is a positive integer that is not a perfect square, and $k \geq 3$ is odd. A natural extension of that work, is to replace $\cot(z)$ with one of the functions $\{\tan(z), \csc(z), \sec(z)\}$. We can settle the tangent and cosecant cases via elementary trigonometric identities:

$$\sum_{n=1}^{\infty} \frac{\tan(\pi n z)}{n^s} = c_s(z) - 2c_s(2z), \quad \sum_{n=1}^{\infty} \frac{\csc(\pi n z)}{n^s} = c_s(z/2) - c_s(z),$$

but it is more challenging to understand the secant zeta function:

$$\psi_s(z) := \sum_{n=1}^{\infty} \frac{\sec(\pi n z)}{n^s}. \quad (1.2)$$

Date: April 5, 2013.

2010 Mathematics Subject Classification. Primary 33E20; Secondary 33B30, 11L03.

Key words and phrases. Secant zeta function, Bernoulli Numbers, Clausen functions, Riemann zeta function.

This work has been partially supported by NSERC Discovery Grant 355412-2008 and FQRNT Subvention établissement de nouveaux chercheurs 144987. The work of FR has also been supported by a Bourse d'été de premier cycle du ISM-CRM.

The main goal of this paper is to prove formulas for special values of $\psi_s(z)$. In Section 2 we prove that the sum converges absolutely if z is an irrational algebraic number and $s \geq 2$. In Section 4 we obtain results such as

$$\psi_2(\sqrt{2}) = -\frac{\pi^2}{3}, \quad \psi_2(\sqrt{6}) = \frac{2\pi^2}{3}.$$

These types of formulas exist because $\psi_k(z)$ obeys a modular transformation which we prove in Section 3 (see equation (3.8)). Furthermore, based on numerical experiments, we conjecture:

Conjecture 1. *Assume that k and j are positive integers, and that k is even. Then $\psi_k(\sqrt{j})\pi^{-k} \in \mathbb{Q}$.*

The results of Section 4 support this conjecture, even though there are still technical hurdles to constructing a complete proof. For instance, we prove that the conjecture holds for infinite subsequences of natural numbers. The rational numbers that appear are also interesting, and we speculate on their properties in the conclusion.

2. CONVERGENCE

Since $\sec(\pi z)$ has poles at the half-integers, it follows that $\psi_s(z)$ is only well-defined if $nz \notin \mathbb{Z} + \frac{1}{2}$ for any integer n . Thus, we exclude rational points with even denominators from the domain of $\psi_s(z)$. If $z = p/q$ with q odd, then $\psi_s(p/q)$ reduces to linear combinations of Hurwitz zeta functions, and (1.2) converges for $s > 1$. Convergence questions become more complicated if z is irrational. Irrationality guarantees that $|\sec(\pi nz)| \neq \infty$, but we still have to account for how often $|\sec(\pi nz)|$ is large compared to n^s . The Thue-Siegel-Roth Theorem gives that $|\sec(\pi nz)| \ll n^{1+\varepsilon}$ when z is algebraic and irrational, and this proves that (1.2) converges for $s > 2$. The case when $s = 2$ requires a more subtle argument. We use a theorem of Worley to show that the set of n 's where $|\sec(\pi nz)|$ is large is sparse enough to ensure that (1.2) converges. We are grateful to Florian Luca for providing this part of the proof. In summary, we have the following theorem:

Theorem 1. *The series in (1.2) converges absolutely in the following cases:*

- (1) *When $z = p/q$ with q odd and $s > 1$.*
- (2) *When z is algebraic irrational, and $s > 2$.*
- (3) *When z is algebraic irrational, and $s = 2$.*

Proof of Theorem 1, parts (1) and (2). Let z be a rational number with odd denominator in reduced form. It is easy to see that the set of real numbers $\{\sec(n\pi z)\}_{n \in \mathbb{N}}$ is finite. Let $M = \max_{n \in \mathbb{N}} |\sec(n\pi z)|$. Then we have

$$\frac{|\sec(\pi nz)|}{n^s} \leq \frac{M}{n^s}.$$

It follows easily from the Weierstrass M -test that (1.2) converges absolutely for $s > 1$.

Now we prove the second part of the theorem. By elementary estimates

$$|\sec(\pi nz)| = |\csc(\pi(nz - 1/2))| \ll \frac{1}{|nz - \frac{1}{2} - k_n|}, \quad (2.1)$$

where k_n is the integer which minimizes $|nz - \frac{1}{2} - k_n|$. Now appeal to the Thue-Siegel-Roth Theorem [Rot55]. In particular, for any algebraic irrational number α , and given $\varepsilon > 0$,

there exists a constant $C(\alpha, \varepsilon)$, such that

$$\left| \alpha - \frac{p}{q} \right| > \frac{C(\alpha, \varepsilon)}{q^{2+\varepsilon}}. \quad (2.2)$$

If we set $\alpha = z$, then (2.1) becomes

$$|\sec(\pi n z)| \ll \frac{1}{n \left| z - \frac{2k_n+1}{2n} \right|} \ll n^{1+\varepsilon}.$$

Therefore we have

$$\frac{|\sec(n\pi z)|}{n^s} \ll \frac{1}{n^{s-1-\varepsilon}},$$

and this implies that (1.2) converges absolutely for $s > 2 + \varepsilon$. Since ε is arbitrarily small the result follows. \square

In order to prove the third part of Theorem 1, we require some background on continued fractions. Recall that any irrational number z can be represented as an infinite continued fraction

$$z = [a_0; a_1, a_2, \dots],$$

and the convergents are given by

$$[a_0; a_1, \dots, a_\ell] = \frac{p_\ell}{q_\ell},$$

which satisfy

$$p_\ell = a_\ell p_{\ell-1} + p_{\ell-2}, \quad (2.3)$$

$$q_\ell = a_\ell q_{\ell-1} + q_{\ell-2}. \quad (2.4)$$

Convergents provide the best possible approximations to algebraic numbers among rational numbers with bounded denominators. In other words, if $0 < q < q_\ell$, then

$$\left| z - \frac{p}{q} \right| > \left| z - \frac{p_\ell}{q_\ell} \right|. \quad (2.5)$$

In addition

$$\frac{1}{q_\ell q_{\ell+1}} > \left| z - \frac{p_\ell}{q_\ell} \right| > \frac{1}{q_\ell (q_{\ell+1} + q_\ell)}. \quad (2.6)$$

Now we state a weak version of a theorem due to Worley [Wor81, Thm. 1].

Theorem 2 (Worley). *Let z be irrational, $k \geq \frac{1}{2}$, and p/q be a rational approximation to z in reduced form for which*

$$\left| z - \frac{p}{q} \right| < \frac{k}{q^2}.$$

Then either p/q is a convergent p_ℓ/q_ℓ to z , or

$$\frac{p}{q} = \frac{ap_\ell + bp_{\ell-1}}{aq_\ell + bq_{\ell-1}}, \quad |a|, |b| < 2k,$$

where a and b are integers.

Now we can complete the proof of Theorem 1. The following proof was kindly provided by Florian Luca.

Proof of Theorem 1, part (3). Let k_n be the integer which minimizes $|nz - \frac{1}{2} - k_n|$. Let W_z denote the set of integers where the quantity is large:

$$W_z = \left\{ n \in \mathbb{N} : \left| nz - \frac{1}{2} - k_n \right| \geq \frac{(\log n)^2}{n} \right\}.$$

Then

$$\sum_{n \in W_z} \frac{|\sec(n\pi z)|}{n^2} \ll |\sec(\pi z)| + \sum_{n=2}^{\infty} \frac{1}{n(\log n)^2},$$

which converges.

Now assume that $n \notin W_z$. Then

$$\left| z - \frac{1 + 2k_n}{2n} \right| < \frac{(\log n)^2}{n^2}.$$

Consider the convergents of z . Let ℓ be such that $q_{\ell-1} \leq 2n < q_\ell$. By Theorem 2 there are at most $O((\log q_\ell)^4)$ solutions to

$$\left| z - \frac{p}{2n} \right| < \frac{(\log n)^2}{n^2} \quad (2.7)$$

with $p \in \mathbb{Z}$ (i.e. consider all values of $|a|, |b| < 2k = 2(\log q_\ell)^2$).

From equations (2.5) and (2.6) we have

$$\left| z - \frac{p}{2n} \right| \geq \left| z - \frac{p_\ell}{q_\ell} \right| \geq \frac{1}{q_\ell(q_{\ell+1} + q_\ell)}.$$

Combining this with equation (2.4) implies

$$\left| nz - \frac{1 + 2k_n}{2} \right| n^2 \geq \frac{n^3}{q_\ell(q_{\ell+1} + q_\ell)} \geq \frac{q_{\ell-1}^3}{8q_\ell(q_\ell(a_{\ell+1} + 1) + q_{\ell-1})} \geq \frac{q_{\ell-1}^3}{8q_\ell^2(a_{\ell+1} + 2)}.$$

Hence, if $n \notin W_z$, we find that

$$\frac{|\sec(n\pi z)|}{n^2} \ll \frac{q_\ell^2(a_{\ell+1} + 2)}{q_{\ell-1}^3} \ll \frac{a_{\ell+1}q_\ell^2}{q_{\ell-1}^3}.$$

Combining the Thue-Siegel-Roth Theorem (equation (2.2)) with (2.6), implies that if z is algebraic

$$\frac{1}{q_\ell q_{\ell+1}} > \left| z - \frac{p_\ell}{q_\ell} \right| > \frac{C(z, \varepsilon)}{q_\ell^{2+\varepsilon}}.$$

Thus $q_{\ell+1} \ll q_\ell^{1+\varepsilon}$. This allows us to place an upper bound on $a_{\ell+1}$:

$$a_{\ell+1} \leq \frac{q_{\ell+1}}{q_\ell} \ll q_\ell^\varepsilon$$

Putting everything together gives the bound

$$\frac{|\sec(n\pi z)|}{n^2} \ll \frac{1}{q_{\ell-1}^{1-4\varepsilon}},$$

and as a result

$$\sum_{n \notin W_z} \frac{|\sec(n\pi z)|}{n^2} \ll \sum_{\ell=1}^{\infty} \frac{(\log q_\ell)^4}{q_\ell^{1-\varepsilon'}} \ll \sum_{\ell=1}^{\infty} \frac{1}{q_\ell^{1-\varepsilon''}}.$$

Since $q_{\ell+1} = a_{\ell+1}q_\ell + q_{\ell-1} \geq q_\ell + q_{\ell-1}$, we conclude that $q_\ell \geq F_\ell$, where F_ℓ denotes the ℓ th Fibonacci number. Since the Fibonacci numbers grow exponentially, we have

$$q_\ell \gg \frac{\varphi^\ell}{\sqrt{5}}, \text{ where } \varphi = \frac{1 + \sqrt{5}}{2}.$$

Setting $\tilde{\varphi} = \varphi^{1-\varepsilon''} > 1$, we finally obtain

$$\sum_{n \notin W_z} \frac{|\sec(n\pi z)|}{n^2} \ll \sum_{\ell=1}^{\infty} \frac{1}{\tilde{\varphi}^\ell} = \frac{1}{\tilde{\varphi} - 1} < \infty.$$

Thus, it follows that (1.2) converges absolutely when $s = 2$. \square

3. A MODULAR TRANSFORMATION FOR $\psi_k(z)$

We begin by noting the following trivial properties of $\psi_s(z)$:

$$\psi_s(-z) = \psi_s(z), \quad (3.1)$$

$$\psi_s(z+2) = \psi_s(z), \quad (3.2)$$

$$2^{1-s}\psi_s(2z) = \psi_s(z) + \psi_s(z+1). \quad (3.3)$$

The main goal of this section is to prove that $\psi_k(z)$ also satisfies a modular transformation. We start from the partial fractions decompositions of $\sec(\pi x)$ and $\csc(\pi x)$, and then perform a convolution trick to obtain an expansion for $\sec(\pi x)\csc(\pi xz)$ (equation (3.5)). Differentiating with respect to x then leads to the transformation for $\psi_k(z)$ (equation (3.8)). The third author used this method to rediscover the Newberger summation rule for Bessel functions [Rog05], [New82]:

$$J_x(y)J_{-x}(y)\frac{\pi}{\sin(\pi x)} = \sum_{n=-\infty}^{\infty} \frac{J_n^2(y)}{n+x}. \quad (3.4)$$

Equation (3.4) follows from applying the convolution trick to partial fractions expansions for $J_x(y)$ and $J_{-x}(y)$ [Lom68].

Lemma 1. *Let $\chi_{-4}(n)$ denote the Legendre symbol modulo 4. Suppose that x and z are selected appropriately.¹ Then*

$$\pi \csc(\pi zx) \sec(\pi x) = \frac{1}{zx} + 8x \sum_{n=1}^{\infty} \frac{\chi_{-4}(n) \csc(\pi n z / 2)}{n^2 - 4x^2} - 2zx \sum_{n=1}^{\infty} \frac{\sec(\pi n(1 + 1/z))}{n^2 - z^2 x^2}. \quad (3.5)$$

Proof. Recall the classical partial fractions expansions [GR94]:

$$\pi (\sec(\pi x) - 1) = 16x^2 \sum_{n=1}^{\infty} \frac{\chi_{-4}(n)}{n(n^2 - 4x^2)}, \quad (3.6)$$

$$\pi \csc(\pi x) - \frac{1}{x} = 2x \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^2 - x^2}. \quad (3.7)$$

¹We assume that z is irrational and algebraic, and x is selected so that the denominators in the sums are never zero.

Both sums converge uniformly. Multiplying the formulas together, expanding via partial fractions, and rearranging the order of summation, we have

$$\begin{aligned}
\pi (\sec(\pi x) - 1) \left(\pi \csc(\pi z x) - \frac{1}{zx} \right) &= 32zx^3 \sum_{\substack{n \geq 1 \\ k \geq 1}} \frac{(-1)^{k+1} \chi_{-4}(n)}{n(n^2 - 4x^2)(k^2 - z^2x^2)} \\
&= 8z^3x^3 \sum_{\substack{n \geq 1 \\ k \geq 1}} \frac{(-1)^{k+1} \chi_{-4}(n)}{n(k^2 - z^2n^2/4)} \left(\frac{1}{z^2n^2/4 - z^2x^2} - \frac{1}{k^2 - z^2x^2} \right) \\
&= 32zx^3 \sum_{n=1}^{\infty} \frac{\chi_{-4}(n)}{n(n^2 - 4x^2)} \left(\sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^2 - z^2n^2/4} \right) \\
&\quad + 32zx^3 \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^2 - z^2x^2} \left(\sum_{n=1}^{\infty} \frac{\chi_{-4}(n)}{n(n^2 - 4k^2/z^2)} \right).
\end{aligned}$$

By (3.6) and (3.7) this becomes

$$\begin{aligned}
\pi (\sec(\pi x) - 1) \left(\pi \csc(\pi z x) - \frac{1}{zx} \right) &= 32x^3 \sum_{n=1}^{\infty} \frac{\chi_{-4}(n)}{n^2(n^2 - 4x^2)} \left(\pi \csc(\pi n z/2) - \frac{2}{nz} \right) \\
&\quad + 2\pi z^3x^3 \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^2(k^2 - z^2x^2)} (\sec(\pi k/z) - 1).
\end{aligned}$$

This reduces to (3.5) after several additional applications of (3.6) and (3.7). We can split up the sums because all of the individual terms converge absolutely. For instance, we can prove that

$$\sum_{k=1}^{\infty} \frac{(-1)^{k+1} \sec(\pi k/z)}{k^2 - z^2x^2},$$

converges absolutely, by showing that the summand is $\ll |\sec(\pi k/z)|k^{-2}$, and then applying Theorem 1. \square

Theorem 3. *Let E_m denote the Euler numbers, and B_m denote the Bernoulli numbers. Suppose that $k \in 2\mathbb{N}$. Then for appropriate choices of z :*

$$\begin{aligned}
(1+z)^{k-1} \psi_k \left(\frac{z}{1+z} \right) - (1-z)^{k-1} \psi_k \left(\frac{z}{1-z} \right) \\
= \frac{(\pi i)^k}{k!} \sum_{m=0}^k (2^{m-1} - 1) B_m E_{k-m} \binom{k}{m} [(1+z)^{m-1} - (1-z)^{m-1}].
\end{aligned} \tag{3.8}$$

Proof. Recall the Taylor series expansions of cosecant and secant:

$$\pi x \csc(\pi x) = - \sum_{m=0}^{\infty} (2^m - 2) B_m \frac{(\pi i x)^m}{m!}, \quad \sec(\pi x) = \sum_{m=0}^{\infty} E_m \frac{(\pi i x)^m}{m!}.$$

Expand both sides of (3.5) in a Taylor series with respect to x . Comparing coefficients yields the following identity for $k \in 2\mathbb{N}$:

$$\frac{(\pi i)^k}{k!} \sum_{m=0}^k (2^{m-1} - 1) B_m E_{k-m} \binom{k}{m} z^{m-1} = -2^k \sum_{n=1}^{\infty} \frac{\chi_{-4}(n) \csc(\pi n z/2)}{n^k} + z^{k-1} \psi_k(1 + 1/z).$$

Finally, let $z \mapsto (1+z)$ and $z \mapsto (1-z)$, and subtract the two results to recover (3.8). The cosecant sums vanish because $\csc(\pi n(1+z)/2) = \csc(\pi n(1-z)/2)$ whenever n is odd. \square

We conclude this subsection with a conjecture on unimodular polynomials. We call a polynomial *unimodular* if its zeros all lie on the unit circle. We have observed numerically that the polynomials in (3.8) have all of their zeros on the vertical line $\operatorname{Re}(z) = 0$. Since the linear fractional transformation $z = (1-x)/(1+x)$ maps the vertical line to the unit circle, we arrive at the following conjecture:

Conjecture 2. *We conjecture that the polynomial*

$$\sum_{m=0}^k 2^m (2^m - 2) B_m E_{k-m} \binom{k}{m} (x - x^m)(1+x)^{k-m}, \quad (3.9)$$

is unimodular when $k \in 2\mathbb{N}$.

This new family of polynomials is closely related to the unimodular polynomials introduced in [LR12], [LS13], and [HZ13].

4. SPECIAL VALUES OF $\psi_k(z)$

Throughout this section we assume that k is a positive even integer. Let $\operatorname{SL}_2(\mathbb{Z})$ denote the set of 2×2 integer-valued matrices with determinant equal to one, and let $\operatorname{PSL}_2(\mathbb{Z}) = \operatorname{SL}_2(\mathbb{Z})/\langle I, -I \rangle$. Recall that $\operatorname{PSL}_2(\mathbb{Z})$ can be identified with the set of linear fractional transformations. The usual group action is

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} z = \frac{az + b}{cz + d}.$$

It is easy to see that multiplying matrices is equivalent to performing compositions of linear fractional transformations. Consider the following matrices in $\operatorname{PSL}_2(\mathbb{Z})$:

$$A = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}.$$

Equations (3.2) and (3.8) are equivalent to

$$\psi_k(Az) = \psi_k(z), \quad (4.1)$$

$$\psi_k(Bz) = (2z+1)^{1-k} \psi_k(z) \quad (4.2)$$

$$+ \frac{(\pi i)^k}{k!} \sum_{m=0}^k (2^{m-1} - 1) B_m E_{k-m} \binom{k}{m} (z+1)^{k-m} \left[(2z+1)^{m-k} - (2z+1)^{1-k} \right].$$

Every matrix $C \in \langle A, B \rangle$ has a factorization of the form $C = A^{j_1} B^{j_2} A^{j_3} \dots$, so equations (4.1) and (4.2) together imply that there exists a z -linear relation between $\psi_k(z)$, $\psi_k(Cz)$, and π^k .

Now we outline a strategy to obtain exact evaluations of $\psi_k(z)$. First select a matrix C , and then find the linear relation between $\psi_k(z)$, $\psi_k(Cz)$, and π^k . Next choose z so that $\psi_k(z) = \psi_k(Cz)$. For example, if $z = 2j + \sqrt{2j(2j+1)}$ in (4.2), then $Bz = z - 4j$

whenever j is a non-zero integer. With some work the equation collapses to

$$\begin{aligned} \psi_k \left(\sqrt{2j(2j+1)} \right) &= \frac{(\pi i)^k}{k!} \sum_{m=0}^k (2^{m-1} - 1) B_m E_{k-m} \binom{k}{m} \\ &\quad \times \left[\frac{\left(1 + \sqrt{\frac{2j}{2j+1}}\right)^{m-1} - \left(1 - \sqrt{\frac{2j}{2j+1}}\right)^{m-1}}{\left(1 + \sqrt{\frac{2j}{2j+1}}\right)^{k-1} - \left(1 - \sqrt{\frac{2j}{2j+1}}\right)^{k-1}} \right], \end{aligned} \quad (4.3)$$

which holds for $j \in \mathbb{Z} \setminus \{0\}$. Similarly, if we take $z = |2j+1| + \sqrt{2j(2j+1)}$, then we arrive at

$$\begin{aligned} \psi_k \left(1 + \sqrt{2j(2j+1)} \right) &= \frac{(\pi i)^k}{k!} \sum_{m=0}^k (2^{m-1} - 1) B_m E_{k-m} \binom{k}{m} \\ &\quad \times \left[\frac{\left(1 + \sqrt{\frac{2j+1}{2j}}\right)^{m-1} - \left(1 - \sqrt{\frac{2j+1}{2j}}\right)^{m-1}}{\left(1 + \sqrt{\frac{2j+1}{2j}}\right)^{k-1} - \left(1 - \sqrt{\frac{2j+1}{2j}}\right)^{k-1}} \right], \end{aligned} \quad (4.4)$$

which also holds for $j \in \mathbb{Z} \setminus \{0\}$. The right-hand sides of (4.3) and (4.4) are invariant under the Galois action $\sqrt{x} \mapsto -\sqrt{x}$, so both formulas are rational with respect to j . Specializing (4.3) at $k=2$ and $k=4$ yields

$$\begin{aligned} \psi_2 \left(\sqrt{2j(2j+1)} \right) &= (3j+1) \frac{\pi^2}{6}, \\ \psi_4 \left(\sqrt{2j(2j+1)} \right) &= \left(\frac{75j^2 + 46j + 6}{8j+3} \right) \frac{\pi^4}{180}. \end{aligned}$$

These results support Conjecture 1. Further evidence for the conjecture is provided by combining (4.3), (4.4), and (3.3), to obtain identities like

$$\psi_2 \left(\sqrt{8j(2j+1)} \right) = \frac{\pi^2}{6}, \quad (4.5)$$

whenever $j \in \mathbb{Z}$.

In general, it seems to be quite difficult to evaluate $\psi_k(\sqrt{j})$ for arbitrary positive integers j . This is due to the fact that (4.1) and (4.2) restrict the available matrices to a subgroup of $\mathrm{PSL}_2(\mathbb{Z})$. If there exists a matrix $C \in \langle A, B \rangle$ which satisfies $C\sqrt{j} = \sqrt{j}$, then we can always evaluate $\psi_k(\sqrt{j})$. We can construct candidate matrices by solving Pell's equation:

$$X^2 - jY^2 = 1.$$

If we choose C to be given by

$$C = \begin{pmatrix} X & jY \\ Y & X \end{pmatrix},$$

then it follows that $C\sqrt{j} = \frac{X\sqrt{j}+jY}{Y\sqrt{j}+X} = \sqrt{j}$, and $\det(C) = X^2 - jY^2 = 1$. Pell's equation has infinitely many integral solutions when $j \geq 1$ [HW38], so there are infinitely many choices of C . The main difficulty is to select appropriate values of X and Y so that C factors into products of A 's and B 's. It is not clear if such a selection is always possible. Notice that $\langle A, B \rangle \subset \Gamma_0(2)$, so it follows $\mathrm{PSL}_2(\mathbb{Z}) \not\subset \langle A, B \rangle$.

We conclude this subsection by noting that $\psi_2(z) = 0$ for infinitely many irrational values of z . Setting $n = -3j$ in Proposition 1 below, yields

$$\psi_2\left(\sqrt{\frac{2(6j^2 - 1)}{3}}\right) = 0, \quad (4.6)$$

for any non-zero integer j . It is worth emphasizing that $\psi_k(z)$ is highly discontinuous with respect to z , so these types of results are not surprising.

Proposition 1. *Suppose that j and n are integers, and $n \neq 0$. Then*

$$\psi_2\left(\sqrt{\frac{2j(2jn + 1)}{n}}\right) = \left(1 + \frac{3j}{n}\right) \frac{\pi^2}{6}. \quad (4.7)$$

Proof. Setting $k = 2$ in (4.2) yields

$$\psi_2(Bz) = \frac{1}{2z + 1} \psi_2(z) + \frac{z(3z^2 + 4z + 2)}{(2z + 1)^2} \frac{\pi^2}{6}. \quad (4.8)$$

Iterating (4.8) gives

$$\psi_2(B^n z) = \frac{1}{2nz + 1} \psi_2(z) + \frac{nz(3z^2 + 4nz + 2)}{(2nz + 1)^2} \frac{\pi^2}{6}. \quad (4.9)$$

The derivation of (4.9) is best accomplished with the aid of a computer algebra system such as *Mathematica*, because significant telescoping occurs on the right. Now consider the matrix

$$C = A^j B^n A^j = \begin{pmatrix} 4jn + 1 & 4j(2jn + 1) \\ 2n & 4jn + 1 \end{pmatrix},$$

and notice that $Cz_0 = z_0$, where

$$z_0 = \sqrt{\frac{2j(2jn + 1)}{n}}.$$

Thus by (4.9) we have

$$\begin{aligned} \psi_2(z_0) &= \psi_2(Cz_0) \\ &= \psi_2(B^n(z_0 + 2j)) \\ &= \frac{1}{2n(z_0 + 2j) + 1} \psi_2(z_0) + \frac{n(z_0 + 2j)(3(z_0 + 2j)^2 + 4n(z_0 + 2j) + 2)}{(2n(z_0 + 2j) + 1)^2} \frac{\pi^2}{6}. \end{aligned}$$

We complete the proof by solving for $\psi_2(z_0)$ and simplifying. \square

5. SPECULATIONS AND CONCLUSION

Assume that k is a positive even integer. Euler gave the following expression for Bernoulli numbers:

$$B_k = -2 \frac{k!}{(2\pi i)^k} \zeta(k). \quad (5.1)$$

Bernoulli numbers are interesting combinatorial objects, so it is natural to ask if the rational part of $\psi_k(\sqrt{j})$ also has interesting properties. This is an obvious question because $\psi_k(0) = \zeta(k)$. For instance, the von Staudt-Clausen theorem gives a complete description of the denominators of Bernoulli numbers:

$$B_k = \sum_{(p-1)|k} \frac{1}{p} + \text{Integer}. \quad (5.2)$$

Is there an analogue of (5.2) for the rational part of $\psi_k(\sqrt{j})$? As an example, consider $\psi_k(\sqrt{6})$ which can be calculated from equation (4.3). In order to eliminate some trivial factors, we define

$$\beta_k := \frac{(3 + \sqrt{6})^{k-1} - (3 - \sqrt{6})^{k-1}}{\sqrt{6}} \frac{k!}{(\pi i)^k} \psi_k(\sqrt{6}).$$

By (4.3) we have

$$\beta_k = \sum_{m=0}^k (2^{m-1} - 1) B_m E_{k-m} \binom{k}{m} 3^{k-m} \frac{(3 + \sqrt{6})^{m-1} - (3 - \sqrt{6})^{m-1}}{\sqrt{6}}.$$

The first few values are $\beta_2 = -8/3$, $\beta_4 = 508/5$, and $\beta_6 = -64896/7$, which all have denominators divisible only by primes where $p - 1 | k$ (as hoped for). The first instance where this fails is β_{20} . The denominator of β_{20} equals $5 \cdot 7 \cdot 11$, but $7 - 1 = 6$ does not divide 20.

Acknowledgements. The authors wish to thank Shabnam Ahktari for helpful discussions and Florian Luca for providing the proof of part (3) of Theorem 1. Finally, the authors are grateful to the referee for comments and suggestions that greatly improved the exposition of this article.

REFERENCES

- [Ber76] B. C. Berndt, Dedekind sums and a paper of G. H. Hardy, *J. London Math. Soc. (2)* **13** (1976), 129–137.
- [GR94] I. S. Gradshteyn and I. M. Ryzhik, Table of Integrals, Series and Products, Academic Press 1994
- [HW38] G. H. Hardy and E. M. Wright, An introduction to the theory of numbers, Clarendon Press (1938).
- [HZ13] H. K. Hwang V. Zacharovas, Limit laws of the coefficients of polynomials with only unit roots, Preprint (2013), arXiv:1301.2021.
- [LR12] M. N. Lalín and M. D. Rogers, Variations of the Ramanujan polynomials and remarks on $\zeta(2j + 1)/\pi^{2j+1}$, to appear in *Funct. Approx. Comment. Math.*.
- [LS13] M. N. Lalín and C. J. Smyth, Unimodularity of roots of self-inversive polynomials, *Acta Math. Hungar.* **138** (2013), no. 1-2, 85–101 .
- [Lew91] L. Lewin, Structural properties of polylogarithms, Mathematical Surveys and Monographs, **37**. American Mathematical Society, Providence, RI, 1991. xviii+412
- [Lom68] E. C. Lommel, Studien uber die Bessel'schen Functionen, Leipzig 1868.
- [New82] B. S. Newberger, New sum rule for products of Bessel functions with application to plasma physics, *J. Math. Phys.* **23** (1982), 1278.
- [Rog05] M. D. Rogers, Partial fractions expansions and identities for products of Bessel functions, *J. Math. Phys.* **46** (2005), 043509.
- [Rot55] K. F. Roth, Rational approximations to algebraic numbers, *Mathematika* **2** (1955), 1–20; corrigendum, 168.
- [Wor81] R. T. Worley, Estimating $\alpha - p/q$. *J. Austral. Math. Soc. Ser. A* **31** (1981), no. 2, 202–206.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MONTREAL, MONTREAL, CANADA
E-mail address: mlalin@dms.umontreal.ca

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MONTREAL, MONTREAL, CANADA
E-mail address: rodriguefrancis@gmail.com

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MONTREAL, MONTREAL, CANADA
E-mail address: mathewrogers@gmail.com